

Isospin-Breaking Vacuum-to- π^0, η Pseudoscalar Matrix Elements at Next-to-Leading Order in the Chiral Expansion

Kim Maltman

*Department of Mathematics and Statistics, York University
4700 Keele St., North York, Ont. CANADA M3J 1P3
Special Research Center for the Subatomic Structure of Matter
University of Adelaide, Australia 5005*

Carl E. Wolfe

*Department of Physics and Astronomy, York University
4700 Keele St., North York, Ont. CANADA M3J 1P3
(February 1, 2008)*

We employ Chiral Perturbation Theory (ChPT) to evaluate the complete set of pseudoscalar matrix elements, $\langle 0|P_f|\pi^0, \eta \rangle$, with P_f any of the flavor-diagonal pseudoscalar currents ($f = u, d, s$), to $\mathcal{O}(m_d - m_u)$, and to next-to-leading order in the chiral expansion. These matrix elements represent the basic input to a QCD sum rule analysis of isospin breaking in the πNN couplings using the three-point-function method. We discuss also how one could use the results to construct a one parameter family of interpolating fields for the π^0 , all of whose members have zero vacuum-to- η matrix element, and explain how this could in principle be used to provide non-trivial tests of the reliability of the assumptions underlying the use of the three-point-function method. It is shown that the isospin-breaking mixing parameters required for this construction receive significant corrections beyond leading order in the chiral expansion.

14.40.Aq, 11.30.Hv, 11.30.Rd, 12.39.Fe

I. INTRODUCTION

QCD sum rules [1] treatments of the isospin-conserving πNN coupling, both using the two-point-function and three-point-function methods [2–8], have proven quite successful. As a consequence, it is natural to consider extending these treatments to an investigation of the strong interaction ($m_d - m_u \neq 0$) contributions to the isospin breaking of these same couplings. The resulting differences among $g_{pp\pi^0}$, $g_{nn\pi^0}$ and $g_{np\pi^+}$ would provide contributions to isospin-breaking observables in few-body systems, and an evaluation based on methods closely related to QCD is, therefore, highly desirable, particularly in light of the recent revival of interest in such phenomena (especially charge symmetry breaking (CSB)) [9–25].

A first attempt in this direction was made in Ref. [25] (MH), where a sum rule analysis using the three-point-function method was performed. In this approach, one considers the three-point functions

$$A_{N'N\pi}(p_1, p_2, q) = \int d^4x_1 d^4x_2 \exp[i(p_1 \cdot x_1 - p_2 \cdot x_2)] \langle 0|T(\eta_{N'}(x_1)P_\pi(0)\bar{\eta}_N(x_2))|0 \rangle \quad (1)$$

where $\eta_{N,N'}$ are (Ioffe) interpolating fields for the nucleons N' , N and P_π a pseudoscalar interpolating field for the pion of interest (which carries momentum $q = p_1 - p_2$). The desired coupling, $g_{N'N\pi}$, then appears as a coefficient in the contribution to the phenomenological (“hadronic”) representation of $A_{N'N\pi}$ having simultaneous poles at $p_1^2 = p_2^2 = M_N^2$ and $q^2 = m_\pi^2$. To evaluate these couplings in terms of vacuum condensates, one then simultaneously evaluates $A_{N'N\pi}$ using the operator product expansion (OPE) (the validity of which requires $q^2 < -1 \text{ GeV}^2$), and relates the hadronic and OPE representations of the correlator via an appropriately Borel transformed dispersion relation [1]. The contributions due to the pion coupling are isolated by keeping only those terms in the OPE proportional to $\not{q}\gamma_5$ (as in the phenomenological representation) which have a $1/q^2$ pole in the limit that the quark masses are neglected [2–4, 25].

It is necessary to exercise some care in the choice of the interpolating field for the neutral pion in Eq. (1). Indeed, if one were to make the most obvious choice, *i.e.*, the $I = 1$ combination

$$P_{\pi_3} = P_{I=1} = P_u - P_d \quad (2)$$

where $P_f = \bar{f}i\gamma_5 f$ is the pseudoscalar current for quark flavor, f , one would find that because, to $\mathcal{O}(m_d - m_u)$, P_{π_3} also couples to the η , the analysis for the extraction of the isospin-breaking πNN couplings was contaminated by η

contributions. This contamination is of the same order in isospin breaking as the πNN isospin-breaking couplings of interest, and hence not negligible [25]. This problem was overcome in MH by the alternate choice

$$P_{\pi^0}^{MH} = (P_u - P_d) + \epsilon^{MH} P_8, \quad (3)$$

where

$$P_8 = \frac{1}{\sqrt{3}} (P_u + P_d - 2P_s) \quad (4)$$

and ϵ^{MH} is to be chosen such that $\langle 0 | P_{\pi^0}^{MH} | \eta \rangle = 0$. To leading order in the chiral expansion, one has, from the results of Gasser and Leutwyler [26],

$$\epsilon^{MH} = \theta_0 = \frac{\sqrt{3}}{4} \left(\frac{m_d - m_u}{m_s - \hat{m}} \right), \quad (5)$$

where θ_0 is the leading order $\pi^0 - \eta$ mixing angle. Since, however, it is known that next-to-leading order contributions to, *e.g.*, f_η are large ($\sim 30\%$), it is likely that similar next-to-leading order corrections to ϵ^{MH} will also be important.

In this paper we will compute the pseudoscalar current matrix elements, $\langle 0 | P_f | \pi^0, \eta \rangle$, required to evaluate these corrections, to $\mathcal{O}(m_d - m_u)$ and to next-to-leading (1-loop) order in the chiral expansion. We will, in addition, show that it is possible to use these matrix elements to construct, not just the field $P_{\pi^0}^{MH}$ above, but a continuous family of similar interpolating fields, all with zero vacuum-to- η matrix element. These fields can then, in principle, be used to study the reliability of the assumptions which go into the use of the three-point-function treatment of the isospin-breaking πNN couplings, a topic we will return to in a later publication.

Let us consider, now, the problem of constructing “pure” π^0 and η interpolating fields from the flavor diagonal pseudoscalar currents. If $\hat{P}_0$ is any $I = 0$ pseudoscalar field combination having non-zero η matrix element, then forming

$$P_{\pi^0} = (P_u - P_d) + \hat{\epsilon}_\pi \hat{P}_0 \quad (6)$$

and requiring $\langle 0 | P_{\pi^0} | \eta \rangle = 0$, *i.e.*,

$$\hat{\epsilon}_\pi = - \frac{\langle 0 | P_u - P_d | \eta \rangle}{\langle 0 | \hat{P}_0 | \eta \rangle}, \quad (7)$$

produces an interpolating field, P_{π^0} , which couples to the physical π^0 , but not at all to the η . A “pure” η interpolating field is constructed analogously:

$$P_\eta = \hat{P}_0 + \hat{\epsilon}_\eta (P_u - P_d) \quad (8)$$

with

$$\hat{\epsilon}_\eta = - \frac{\langle 0 | \hat{P}_0 | \pi^0 \rangle}{\langle 0 | P_u - P_d | \pi^0 \rangle}. \quad (9)$$

The basic ingredients in the construction of P_{π^0} , P_η are, therefore, the vacuum-to- π^0, η matrix elements of the $I = 1$ ($P_{I=1}$) and $I = 0$ ($P_u + P_d$ and P_s) pseudoscalar currents, where π^0, η refer to the physical (isospin-mixed) pseudoscalar mesons. It is these matrix elements that we will evaluate to $\mathcal{O}(m_d - m_u)$, and to 1-loop order in Chiral Perturbation Theory (ChPT) (next-to-leading order in the chiral expansion) below.

II. EVALUATION OF THE VACUUM-TO- π^0, η MATRIX ELEMENTS OF THE PSEUDOSCALAR CURRENTS

Having explained above why an evaluation of the isospin-breaking pseudoscalar current vacuum-to- π^0, η matrix elements is of interest, we now turn to the explicit evaluation of these quantities, for which we employ the methods of ChPT [26]. In order to obtain both the leading and next-to-leading contributions in the chiral expansion of these quantities, it is necessary to include those pieces of the effective Lagrangian up to and including fourth order in the chiral counting [27], in the presence of scalar and pseudoscalar external sources, s and p . The relevant terms, as worked out by Gasser and Leutwyler [26], are

$$\mathcal{L}_{eff} = \mathcal{L}^{(2)} + \mathcal{L}^{(4)} \quad (10)$$

where

$$\mathcal{L}^{(2)} = \frac{1}{4}f^2\text{Tr}(\partial_\mu U \partial^\mu U^\dagger) + \frac{1}{4}f^2\text{Tr}[\chi^\dagger U + \chi U^\dagger] \quad (11)$$

and

$$\begin{aligned} \mathcal{L}^{(4)} = & L_1 [\text{Tr}(\partial_\mu U \partial^\mu U^\dagger)]^2 + L_2 \text{Tr}(\partial_\mu U \partial_\nu U^\dagger) \text{Tr}(\partial^\mu U \partial^\nu U^\dagger) + L_3 \text{Tr}(\partial_\mu U^\dagger \partial^\mu U \partial_\nu U^\dagger \partial^\nu U) \\ & + L_4 \text{Tr}(\partial_\mu U \partial^\mu U^\dagger) \text{Tr}(\chi^\dagger U + \chi U^\dagger) + L_5 \text{Tr}[\partial_\mu U^\dagger \partial^\mu U (\chi^\dagger U + U^\dagger \chi)] \\ & + L_6 [\text{Tr}(\chi^\dagger U + \chi U^\dagger)]^2 + L_7 [\text{Tr}(\chi^\dagger U - \chi U^\dagger)]^2 + L_8 \text{Tr}(\chi^\dagger U \chi^\dagger U + \chi U^\dagger \chi U^\dagger) \\ & + H_2 \text{Tr}(\chi^\dagger \chi) \end{aligned} \quad (12)$$

where $U = \exp(i\vec{\lambda}\cdot\vec{\pi}/f)$, with $\{\pi^a\}$ the octet of pseudoscalar Goldstone boson fields and $\{\lambda^a\}$ the usual Gell-Mann matrices, f is a (second order) low-energy constant (LEC) which turns out to be equal to the pion decay constant in the chiral limit,

$$\chi = 2B_0(s + ip) \quad (13)$$

with B_0 another second order LEC, related to the quark condensate in the chiral limit, and $\{L_k\}$, H_2 are fourth order LEC's. Terms in the general fourth order piece of the effective Lagrangian, $\mathcal{L}_{eff}$, which vanish in the absence of external vector and axial vector sources have been dropped in writing Eqs. (11), (12). The second and fourth order LEC's appearing in these equations are not determined by the symmetry arguments which go into constructing $\mathcal{L}_{eff}$ and must be determined phenomenologically. (See, for example, Ref. [28] for a recent up-to-date tabulation). For our purposes, the external scalar source is to be set equal to the quark mass matrix, M , in order to correctly incorporate the effects of explicit chiral symmetry breaking, and the external pseudoscalar source is to be retained to first order, in order to generate the correct representation of the underlying QCD pseudoscalar currents in the low-energy effective theory.

From the form of $\mathcal{L}_{eff}$ given above, it is straightforward to obtain the following low-energy representations of the pseudoscalar currents in terms of the Goldston boson fields $\{\pi^a\}$ occuring in U :

$$\begin{aligned} P_u = & B_0 f \left[\left(\pi_3 + \frac{\pi_8}{\sqrt{3}} \right) - \frac{1}{6f^2} \left(\pi_3^3 + \sqrt{3}\pi_3^2\pi_8 + \pi_3\pi_8^2 + \frac{\pi_8^3}{3\sqrt{3}} + 2\pi^+\pi^-\pi_3 \right. \right. \\ & \left. \left. + \frac{6}{\sqrt{3}}\pi^+\pi^-\pi_8 + 4K^+K^-\pi_3 + 4K^0\bar{K}^0\pi_3 + 2\sqrt{2}\pi^+K^-K^0 + 2\sqrt{2}\pi^-K^+\bar{K}^0 \right) \right] \\ & + \frac{32B_0^2}{f} \left[(m_s + 2\hat{m}) \left(\pi_3 + \frac{\pi_8}{\sqrt{3}} \right) L_6 + \left((m_u - m_d)\pi_3 + \frac{2}{\sqrt{3}}(\hat{m} - m_s)\pi_8 \right) L_7 \right. \\ & \left. + m_u \left(\pi_3 + \frac{\pi_8}{\sqrt{3}} \right) L_8 \right] + \dots \\ P_d = & B_0 f \left[\left(-\pi_3 + \frac{\pi_8}{\sqrt{3}} \right) - \frac{1}{6f^2} \left(-\pi_3^3 + \sqrt{3}\pi_3^2\pi_8 - \pi_3\pi_8^2 + \frac{\pi_8^3}{3\sqrt{3}} - 2\pi^+\pi^-\pi_3 \right. \right. \\ & \left. \left. + \frac{6}{\sqrt{3}}\pi^+\pi^-\pi_8 - 4K^+K^-\pi_3 - 4K^0\bar{K}^0\pi_3 + 2\sqrt{2}\pi^+K^-K^0 + 2\sqrt{2}\pi^-K^+\bar{K}^0 \right) \right] \\ & + \frac{32B_0^2}{f} \left[(m_s + 2\hat{m}) \left(-\pi_3 + \frac{\pi_8}{\sqrt{3}} \right) L_6 + \left((m_u - m_d)\pi_3 + \frac{2}{\sqrt{3}}(\hat{m} - m_s)\pi_8 \right) L_7 \right. \\ & \left. + m_d \left(-\pi_3 + \frac{\pi_8}{\sqrt{3}} \right) L_8 \right] + \dots \\ P_s = & B_0 f \left[-\frac{2}{\sqrt{3}}\pi_8 - \frac{1}{6f^2} \left(-8\frac{\pi_8^3}{3\sqrt{3}} + 2K^+K^-\pi_3 - 2K^0\bar{K}^0\pi_3 - 2K^0\bar{K}^0\pi_3 - \frac{6}{\sqrt{3}}K^+K^-\pi_8 \right. \right. \end{aligned}$$

$$\begin{aligned}
& -\frac{6}{\sqrt{3}}K^0\bar{K}^0\pi_8 + 2\sqrt{2}\pi^+K^-K^0 + 2\sqrt{2}\pi^-K^+\bar{K}^0 \Big) \Big] + \frac{32B_0^2}{f} \Big[-\frac{2}{\sqrt{3}}(m_s + 2\hat{m})\pi_8L_6 \\
& + \left((m_u - m_d)\pi_3 + \frac{2}{\sqrt{3}}(\hat{m} - m_s)\pi_8 \right) L_7 - \frac{2}{\sqrt{3}}m_s\pi_8L_8 \Big] + \dots
\end{aligned} \tag{14}$$

where π_3 and π_8 are the pure $I = 1, 0$ unrenormalized octet fields appearing in the matrix variable U , $\hat{m} = (m_u + m_d)/2$, and we have written down explicitly only those terms which will be required for our calculations below.

Using these representations, it is then straightforward to compute the matrix elements in question. The set of graphs which contribute is shown in Fig. 1. For practical purposes it is convenient to recast the low-energy representation of the currents in terms of the renormalized, physical π^0 , η fields, rather than unrenormalized octet fields π_3 and π_8 . In order to correctly incorporate all effects at next-to-leading order in the chiral expansion, it is necessary, for those terms linear in π_3 , π_8 and not proportional to the fourth order LEC's $L_{6,7,8}$ in the representations of the currents, to use a transformation between the π_3 , π_8 and π^0 , η bases which is valid to 1-loop order. The relevant expressions are [16]:

$$\begin{aligned}
\pi^0 &= Z_3^{-1/2} (\pi_3 + \hat{\theta}_1 \pi_8) \\
\eta &= Z_8^{-1/2} (-\hat{\theta}_2 \pi_3 + \pi_8)
\end{aligned} \tag{15}$$

where Z_3 and Z_8 are the π_3 and π_8 wavefunction renormalization constants to 1-loop order in the isospin symmetry limit and

$$\begin{aligned}
\hat{\theta}_1 &= \theta_0 \left[1 - \frac{7}{3}\mu_\pi + \frac{4}{3}\mu_K + \mu_\eta + \left(\frac{3m_\eta^2 + m_\pi^2}{64\pi^2 f^2} \right) \left(1 + \frac{m_\pi^2}{\bar{m}_K^2 - m_\pi^2} \log(m_\pi^2/\bar{m}_K^2) \right) \right. \\
&\quad \left. + \left(\frac{m_\eta^2 - m_\pi^2}{64\pi^2 f^2} \right) (1 + \log(\bar{m}_K^2/\mu^2)) - \frac{32(\bar{m}_K^2 - m_\pi^2)}{f^2} (3L_7^r + L_8^r) \right] \\
\hat{\theta}_2 &= \theta_0 \left[1 - \frac{11}{3}\mu_\pi + \frac{8}{3}\mu_K + \mu_\eta + \left(\frac{3m_\eta^2 + m_\pi^2}{64\pi^2 f^2} \right) \left(1 + \frac{m_\pi^2}{\bar{m}_K^2 - m_\pi^2} \log(m_\pi^2/\bar{m}_K^2) \right) \right. \\
&\quad \left. - \left(\frac{m_\eta^2 - m_\pi^2}{64\pi^2 f^2} \right) (1 + \log(\bar{m}_K^2/\mu^2)) - \frac{32(\bar{m}_K^2 - m_\pi^2)}{F^2} (3L_7^r + L_8^r) \right].
\end{aligned} \tag{16}$$

In Eqs. (16), $\mu_P \equiv [m_P^2 \log(m_P^2/\mu^2)]/32\pi^2 f^2$, with μ the (dimensional regularization) renormalization scale, $L_k^r(\mu)$ are the renormalized version of the fourth order LEC's, and $\bar{m}_K^2$ is the average of the neutral and charged kaon squared masses. Standard manipulations then lead to the following expressions for the vacuum-to- π^0 , η matrix elements, valid to $\mathcal{O}(m_d - m_u)$, and to fourth order in the chiral expansion:

$$\langle 0 | P_u - P_d | \pi^0 \rangle = \frac{m_\pi^2 f_\pi}{\hat{m}} \tag{17}$$

$$\begin{aligned}
\langle 0 | P_u + P_d | \pi^0 \rangle &= \frac{m_\pi^2 f_\pi \theta_0}{\sqrt{3}\hat{m}} \left[1 + \left(\frac{-7\mu_\pi + 4\mu_K + 3\mu_\eta}{f^2} \right) + \left(\frac{3m_\eta^2 + m_\pi^2}{64\pi^2 f^2} \right) \left(1 + \right. \right. \\
&\quad \left. \left[\frac{m_\pi^2}{\bar{m}_K^2 - m_\pi^2} \right] \log \left(\frac{m_\pi^2}{\bar{m}_K^2} \right) \right) + \frac{(\bar{m}_K^2 - m_\pi^2)}{16\pi^2 f^2} (1 + \log(m_K^2/\mu^2)) \\
&\quad \left. - \frac{288B_0}{f^2} (m_s - \hat{m})L_7^r - \frac{96B_0}{f^2} (m_s - \hat{m})L_8^r \right]
\end{aligned} \tag{18}$$

$$\begin{aligned}
\langle 0 | P_s | \pi^0 \rangle &= -\frac{m_\pi^2 f_\pi \theta_0}{\sqrt{3}\hat{m}} \left[1 + \left(\frac{-2\mu_\pi + 2\mu_K}{f^2} \right) - \frac{B_0(m_s - \hat{m})}{16\pi^2 f^2} (1 + \log(m_K^2/\mu^2)) \right. \\
&\quad \left. + \frac{B_0(m_s + \hat{m})}{16\pi^2 f^2} \left(1 + \frac{m_\pi^2}{(\bar{m}_K^2 - m_\pi^2)} \log \left(\frac{m_\pi^2}{\bar{m}_K^2} \right) \right) \right]
\end{aligned} \tag{19}$$

$$\begin{aligned}
\langle 0|P_u - P_d|\eta \rangle = & -\frac{m_\pi^2 f_\pi \theta_0}{\hat{m}} \left[1 + \left(\frac{-7\mu_\pi + 6\mu_K + \mu_\eta}{3f^2} \right) + \left(\frac{3m_\eta^2 + m_\pi^2}{64\pi^2 f^2} \right) \left(1 + \right. \right. \\
& \left. \left[\frac{m_\pi^2}{m_K^2 - m_\pi^2} \right] \log \left(\frac{m_\pi^2}{m_K^2} \right) \right) + \frac{(m_\eta^2 - m_\pi^2)}{64\pi^2 f^2} (1 + \log(m_K^2/\mu^2)) \\
& \left. - \frac{16B_0}{3f^2}(m_s - \hat{m})L_5^r - \frac{96B_0}{f^2}(m_s - \hat{m})L_7^r - \frac{32B_0}{f^2}(m_s - \hat{m})L_8^r \right] \quad (20)
\end{aligned}$$

$$\langle 0|P_u + P_d|\eta \rangle = \frac{m_\pi^2 f_\pi}{\sqrt{3}\hat{m}} \left[1 + \left(\frac{-2\mu_\pi + 2\mu_K}{f^2} \right) - \frac{16B_0}{3f^2}(m_s - \hat{m})L_5^r - \frac{64B_0}{f^2}(m_s - \hat{m})L_7^r \right] \quad (21)$$

$$\begin{aligned}
\langle 0|P_s|\eta \rangle = & -\frac{m_\pi^2 f_\pi}{\sqrt{3}\hat{m}} \left[1 + \left(\frac{\mu_\pi - \mu_\eta}{f^2} \right) - \frac{16B_0}{3f^2}(m_s - \hat{m})L_5^r + \frac{32B_0}{f^2}(m_s - \hat{m})L_7^r \right. \\
& \left. + \frac{32B_0}{f^2}(m_s - \hat{m})L_8^r \right]. \quad (22)
\end{aligned}$$

The cancellation of divergences, and the overall scale-independence of the expressions for the matrix elements above, provide non-trivial checks on the calculations, as do the relations

$$\langle 0|\partial_\mu A_f^\mu|P \rangle = \langle 0|2m_f P_f|P \rangle + \text{anomalous terms} \quad (23)$$

which follow from the Ward identities between the divergence of the axial currents and the pseudoscalar currents. (Note that the isospin-breaking vacuum-to- π^0, η matrix elements of the axial currents A_8^μ and A_3^μ have been evaluated previously by Gasser and Leutwyler [26], and that the anomalous terms in Eq. (23) cancel when one considers the combinations which enter A_3^μ and A_8^μ . The vacuum-to-pseudoscalar matrix elements of the anomalous terms are given explicitly in Ref. [29].) In bringing the expressions for the matrix elements into the forms displayed above, we have made use of the 1-loop expressions for m_π^2 and f_π [26]. The appearance of the common multiplicative factor $m_\pi^2 f_\pi / \hat{m}$ in all the matrix elements has no physical significance; the equations are simply written in this form in order to simplify the later calculation of the mixing parameters relevant to the “pure” π^0 interpolating field.

III. INTERPOLATING FIELDS FOR THE NEUTRAL PION

As explained above, the MH π^0 interpolating field choice Eq. (3) is not unique. In fact, if we consider

$$P(\theta) = (P_u - P_d) + \epsilon(\theta) (\cos \theta P_8 + \sin \theta P_0) \quad (24)$$

where $P_0 = (P_u + P_d + P_s)/\sqrt{3}$, the results of Eqs. (18), (19) show that, in general,

$$\langle 0|\cos \theta P_8 + \sin \theta P_0|\eta \rangle \neq 0. \quad (25)$$

As a result, one can, using the construction above, chose $\epsilon(\theta)$ in Eq. (24) in such a way that $\langle 0|P(\theta)|\eta \rangle = 0$ for any value of θ . Any such $P(\theta)$ can then be employed in an analogue of the MH analysis of the three point correlator, in the sense that, in all such analyses, the unwanted η contributions will have been removed by the corresponding choice of $\epsilon(\theta)$.

The existence of the family of π^0 interpolating fields above is potentially useful as a means of investigating features of the three-point-function method that have been the subject of criticism in the literature. The most serious such criticism of the method as a way of estimating the πNN couplings is the objection that it requires making the assumption of pion dominance in the pseudoscalar channel all the way out to values of q^2 of order -1 GeV^2 for which one can begin to hope that the operator product expansion (OPE) analysis of the correlator might be valid [8]. The danger, of course, is that, so far from the pion pole, the contributions of other resonances (η' , $\eta(1295)$, $\pi(1300)$, etc.) might no longer be negligible. Although MH have not attempted to investigate this aspect of the method in detail, they argue that their choice of π^0 interpolating field (corresponding to $\theta = 0$, and the leading chiral order estimate for $\epsilon(\theta)$ above) represents the unique one that, at least to leading order, also has no overlap with the η' . At first sight this seems a plausible argument since, in the $SU(3)_F$ and isospin symmetry limits, $P_u - P_d$ couples only to $I = 1$ states and P_8 only to flavor octet states. On closer inspection, however, one can see that $\langle 0|P(0)|\eta' \rangle \neq 0$ [30],

so that there remain potentially dangerous higher resonance contaminations. Although this point is part of a more detailed analysis of higher resonance contaminations in the three-point-function approach, the results of which will be presented elsewhere, we will briefly outline the key aspects of the argument below. The main point of relevance to the present discussion is that, although one can, in principle, find a value for θ for which at least the vacuum-to- η' matrix element vanishes, the information required to actually determine this value is not available. Thus, for any particular choice of θ , one has a coupling of $P(\theta)$ to the η' of unknown strength (in addition to potential couplings to other higher pseudoscalar resonances), which presents a serious problem for the three-point-function method.

One way of dealing with this problem is to investigate the extracted values of the isospin-breaking πNN couplings as a function of θ . If one finds a region of θ in which the results are not sensitive to variations of θ , then one will have some *a posteriori* justification for assuming that the channel of interest is, indeed, dominated by the lowest-lying (π^0) state. Note that the coupling of $P(\theta)$ to the physical π^0 , from Eqs. (17)–(19) above, is, to $\mathcal{O}(m_d - m_u)$, $m_\pi^2 f_\pi^2 / \hat{m}$, independent of θ . The vacuum-to- π^0, η matrix elements above provide the essential input to such an analysis, and we will recast them in a form suitable for use in such an analysis below.

Before doing so, however, let us briefly outline why the MH field choice, $\theta = 0$, does not actually remove the coupling to the η' . Recall that the quark mass matrix responsible for flavor and isospin breaking can be decomposed as

$$M = -\frac{1}{2}(m_d - m_u)\lambda_3 - \frac{1}{\sqrt{3}}(m_s - \hat{m})\lambda_8 + \frac{1}{3}(m_s + 2\hat{m}) . \quad (26)$$

To leading order in the isospin-breaking and flavor-breaking mass differences, therefore, the vacuum-to- η' matrix elements of $P_u - P_d$ and P_8 are related simply by the ratio of mass-dependent coefficients of λ_3 and λ_8 in Eq. (26). It then follows straightforwardly [30] that, to leading order in the masses,

$$\langle 0|P(0)|\eta' \rangle = 3\theta_0 \langle 0|P_8|\eta' \rangle \quad (27)$$

where $\langle 0|P_8|\eta' \rangle = \mathcal{O}(m_s - \hat{m})$. The RHS is thus non-zero, and $\mathcal{O}(m_d - m_u)$. If one considers $\theta \neq 0$, of course, one has, in addition to the $8_F \times 8_F \rightarrow 1_F$ reduced matrix element which governs $\langle 0|P_u - P_d|\eta' \rangle$ and $\langle 0|P_8|\eta' \rangle$, the $1_F \times 1_F \rightarrow 1_F$ reduced matrix element relevant to $\langle 0|P_0|\eta' \rangle$. One can then certainly, in principle, find a θ such that $\langle 0|P(\theta)|\eta' \rangle = 0$. The problem is that, even to do so at leading order in the quark masses, one would need to know the ratio of the two reduced matrix elements above, and this information is not available. Moreover, even if it were, this would not necessarily ensure that, for such a value of θ , the couplings of the higher resonances were small for the same value of θ . We will return to these issues, and to a discussion of the reliability of the three-point-function method, in more detail in a future publication [30].

Let us return, then, to our 1-loop evaluation of $\epsilon(\theta)$. From Eqs. (7), (20), (21) and (22) it follows that

$$\epsilon(\theta) = \left[\frac{\theta_0}{\cos \theta} \right] \frac{N}{D} \quad (28)$$

where

$$\begin{aligned} N = & \left[1 + \left(\frac{-7\mu_\pi + 6\mu_K + \mu_\eta}{3f^2} \right) + \left(\frac{3m_\eta^2 + m_\pi^2}{64\pi^2 f^2} \right) \left(1 + \left[\frac{m_\pi^2}{\bar{m}_K^2 - m_\pi^2} \right] \log \left(\frac{m_\pi^2}{m_K^2} \right) \right) \right. \\ & + \frac{(m_\eta^2 - m_\pi^2)}{64\pi^2 f^2} (1 + \log(m_K^2/\mu^2)) - \frac{16B_0}{3f^2} (m_s - \hat{m})L_5^r \\ & \left. - \frac{96B_0}{f^2} (m_s - \hat{m})L_7^r - \frac{32B_0}{f^2} (m_s - \hat{m})L_8^r \right] \end{aligned} \quad (29)$$

$$\begin{aligned} D = & \left[1 + \left(\frac{-3 \tan \theta \mu_\pi + 2(1 + \tan \theta)\mu_K + (\tan \theta - 2)\mu_\eta}{3f^2} \right) - \frac{16B_0(m_s - \hat{m})}{3f^2} L_5^r \right. \\ & \left. - \frac{32B_0(m_s - \hat{m})}{f^2} \tan \theta L_7^r - \frac{32B_0(m_s - \hat{m})}{3f^2} (\tan \theta - 2)L_8^r \right] . \end{aligned} \quad (30)$$

The factor $\theta_0/\cos \theta$ represents the leading order contribution to $\epsilon(\theta)$ in the chiral expansion and, of course, reproduces ϵ^{MH} (corresponding to $\theta = 0$) at this order. The remaining factors represent the next-to-leading order corrections. We will discuss their magnitude below. One should note that, as $\theta \rightarrow \pi/2, 3\pi/2$, the denominator of Eq. (28) tends to a finite value involving the chiral log and fourth order LEC terms. This means that, near such values of θ , $\epsilon(\theta)$ becomes

large, and with significant uncertainties (associated with uncalculated next-to-next-to-leading order contributions to the denominator). As such, the corresponding interpolating field choices, though in principle reasonable, are not useful practically. In order to avoid these regions of θ it is convenient to consider a re-parametrization of the family of interpolating fields in which one writes

$$\begin{aligned} P(\alpha) &= (P_u - P_d) + \epsilon(\alpha) \left[\alpha(P_u + P_d) + (1 + \alpha)P_s \right] \\ &= (P_u - P_d) + \epsilon(\alpha) \frac{1}{\sqrt{3}} \left[-P_8 + (3\alpha + 1)P_0 \right], \end{aligned} \quad (31)$$

where, as usual, $\epsilon(\alpha)$ is to be chosen for each α in such a way that

$$\langle 0 | P(\alpha) | \eta \rangle = 0. \quad (32)$$

The second line of Eq. (31) shows that such a form for $P(\alpha)$ represents a simple reparametrization of $P(\theta)$, in which a factor of $-\sqrt{3}\cos\theta$ has been absorbed into ϵ and $\tan\theta$ has been rewritten as $3\alpha + 1$. The regions of θ we wish to avoid then correspond to $\alpha \rightarrow \infty$. One then easily shows that Eq. (32) implies

$$\begin{aligned} \epsilon(\alpha) &= -\sqrt{3}\theta_0 \left[1 + \left(\frac{-(10 + 9\alpha)\mu_\pi + 6(1 + \alpha)\mu_K + (4 + 3\alpha)\mu_\eta}{3f^2} \right) \right. \\ &\quad \left. - \frac{32}{3f^2} (4 + 3\alpha) (\bar{m}_K^2 - m_\pi^2) (3L_7^r + L_8^r) + \left(\frac{3m_\eta^2 + m_\pi^2}{64\pi^2 f^2} \right) \left(1 + \right. \right. \\ &\quad \left. \left. \left[\frac{m_\pi^2}{\bar{m}_K^2 - m_\pi^2} \right] \log \left(\frac{m_\pi^2}{\bar{m}_K^2} \right) \right) + \frac{(m_\eta^2 - m_\pi^2)}{64\pi^2 f^2} (1 + \log(m_K^2/\mu^2)) \right] \\ &= -\sqrt{3}\theta_0 [1.299 + 0.0894\alpha], \end{aligned} \quad (33)$$


where, in arriving at Eq. (33), we have made an expansion of the denominator in Eq. (7) so that the whole expression is now explicitly consistent to 1-loop order. The reparametrization has been chosen such that, at leading order in the chiral expansion, $\epsilon(\alpha)$ reduces to $-\sqrt{3}\theta_0$, so the leading 1 in the square brackets in Eq. (33) represents the tree-level contribution. The remaining terms are the 1-loop corrections and, as expected, are not small. This can be seen from the last line of Eq. (33), where the deviation of the term $1.299 + 0.0894\alpha$ from 1 given the size of the 1-loop corrections. For reference sake we note that the MH choice $\theta = 0$ corresponds to $\alpha = -1/3$, and the choice in which there is no P_s content to the interpolating field corresponds to $\alpha = -1$. (To completely convert back to θ notation for the MH case, one must also absorb the constant $-1/\sqrt{3}$ back into the definition of ϵ .) The enhancements due to next-to-leading order corrections are, in these cases, 1.27 and 1.21, respectively. Amusingly, the choice $\alpha = -4/3$ removes the LEC contributions completely from the expression for ϵ .

IV. SUMMARY

We have evaluated the leading $\mathcal{O}(m_d - m_u)$ strong isospin-breaking contributions to the matrix elements $\langle 0 | P_f | \pi^0, \eta \rangle$ at next-to-leading order in the chiral expansion. The results are given in Eqs. (17)–(22). We have also showed how to use this information to construct a family of π^0 interpolating fields which could, in future, be employed to investigate the reliability of analyses which attempt to extract the isospin-breaking πNN couplings via the three point function method in QCD sum rules. The isospin-breaking parameters, $\epsilon(\alpha)$, required in this construction in order to remove unwanted η contaminations from the extracted π^0 couplings, were shown to receive significant contributions at next-to-leading order. As a result, if one wishes to remove such contamination, it is essential that the full next-to-leading order expressions, given in Eq. (33), be used.

ACKNOWLEDGMENTS

The authors acknowledge the continuing financial support of the Natural Sciences and Engineering Research Council of Canada. KM would also like to acknowledge useful conversations with Terry Goldman, Derek Leinweber, Tony Williams and Mike Birse, as well as the hospitality of the Special Research Center for the Subatomic Structure of Matter at the University of Adelaide, where much of this work was performed.



vacpifig1.ps

FIG. 1. Graphs contributing to the vacuum-to- π^0 , η matrix elements. The wavy lines represent the pseudoscalar currents, the solid dots the vertices generated by the $\mathcal{O}(q^2)$ part $\mathcal{L}_{eff}$, the ‘X’ and the circle enclosing an ‘x’ vertices generated by the $\mathcal{O}(q^4)$ part of $\mathcal{L}_{eff}$.

-
- [1] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, Nucl. Phys. **B147**, 385, 448 (1979).
 - [2] L.J. Reinders, H. Rubinstein and S. Yazaki, Nucl. Phys. **B213**, 109 (1983).
 - [3] L.J. Reinders, Act. Phys. Pol. **15**, 329 (1984).
 - [4] L.J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rep. **127**, 1 (1985).
 - [5] S. Narison and N. Paver, Phys. Lett. **B135**, 159 (1984).
 - [6] S. Narison, *QCD Spectral Sum Rules* (World Scientific, Singapore, 1989).
 - [7] H. Shiomi and T. Hatsuda, Nucl. Phys. **A594**, 294 (1995).
 - [8] M. Birse and B. Krippa, Phys. Lett. **B373**, 9 (1996).
 - [9] G.A. Miller, B.M.K. Nefkens and I. Slaus, Phys. Rep. **194**, 1 (1990).
 - [10] T. Goldman, J. A. Henderson and A. W. Thomas, Few-Body Systems **12**,123 (1992).
 - [11] J. Piekarewicz and A. G. Williams, Phys. Rev. **C47**, R2461 (1993); J. Piekarewicz, Phys. Rev. **C48**, 1555 (1993), Phys. Lett. **B358**, 27 (1995).
 - [12] G. Krein, A. W. Thomas, and A. G. Williams, Phys. Lett. **B317**, 293 (1993); H. O’Connell, B. Pearce, A.W. Thomas and A.G. Williams, Phys. Lett **B336**, 1 (1994); H. O’Connell, B. Pearce, A.W. Thomas and A.G. Williams, Phys. Lett. **B354**, 14 (1995); H. O’Connell, A.G. Williams, M. Bracco and G. Krein, Phys. Lett. **B370**, 12 (1996).
 - [13] K. Maltman and T. Goldman, Nucl. Phys. **A572**, 682 (1994).

- [14] C.-T. Chan, E.M. Henley and T. Meissner, Phys. Lett. **B343**, 7 (1995).
- [15] T. Hatsuda, E.M. Henley, T. Meissner and G. Krein, Phys. Rev. **C49**, 452 (1994).
- [16] K. Maltman, Phys. Lett. **B313**, 203 (1993); Phys. Lett. **B351**, 56 (1995).
- [17] K. Maltman, Phys. Rev. **D53**, 2563, 2573 (1996); Phys. Lett. **B362**, 11 (1995).
- [18] K. L. Mitchell, P. C. Tandy, C. D. Roberts and R. T. Cahill, Phys. Lett. **B335** (1994) 282; M.R. Frank and P.C. Tandy, Phys. Rev. **C49**, 478 (1994); M.R. Frank, Phys. Rev. **C51**, 987 (1995); K.L. Mitchell and P.C. Tandy, Phys. Rev. **C55**, 1477 (1997).
- [19] R. Urech, Phys. Lett. **B355**, 308 (1995); H. Neufeld and H. Rupertsberger, Z. Phys. **C68**, 91 (1995).
- [20] S. Gardner, C.J. Horowitz and J. Piekarewicz, Phys. Rev. Lett. **75**, 2462 (1995); Phys. Rev. **C53**, 1143 (1996).
- [21] U. van Kolck, J. Friar and T. Goldman, Phys. Lett. **B371**, 169 (1996).
- [22] M.J. Iqbal, X.-M. Jin and D.B. Leinweber, Phys. Lett. **B367**, 45 (1996); Phys. Lett. **B386**, 55 (1996).
- [23] T.D. Cohen and G.A. Miller, Phys. Rev. **C52**, 3428 (1995).
- [24] K. Maltman, H. O'Connell and A.G. Williams, Phys. Lett. **B376**, 19 (1996).
- [25] T. Meissner and E.M. Henley, nucl-th/9601030, Phys. Rev. **C**, in press.
- [26] J. Gasser and H. Leutwyler, Nucl. Phys. **B250** (1985) 465
- [27] S. Weinberg, Physica **96A**, 327 (1979).
- [28] V. Bernard, N. Kaiser and U.-G. Meissner, Int. J. Mod. Phys. **E4**, 193 (1995).
- [29] J.F. Donoghue and D. Wyler, Phys. Rev. **D45**, 892 (1992).
- [30] K. Maltman, "Higher Resonance Contamination of πNN Couplings Obtained Via the Three-Point Function Method in QCD Sum Rules", York University preprint, June 1997.

